

OXIDATIVE STRESS IN TYPE 2 DIABETES: MECHANISMS AND THE POTENTIAL OF ANTIOXIDANT THERAPY

Farangiz Sunnatova Akbarovna

3rd year student of Samarkand state medical university

Abstract: Type 2 diabetes mellitus (T2DM) is one of the most common chronic diseases worldwide. Alongside insulin resistance and impaired glucose metabolism, oxidative stress plays a central role in its development and complications. Oxidative stress arises from excessive production of reactive oxygen species (ROS) and reduced antioxidant defense, leading to cellular damage. This article reviews the biochemical mechanisms of oxidative stress in T2DM, its harmful effects on lipids, proteins, and DNA, and the role of antioxidants as potential therapeutic agents. Vitamins C and E, glutathione, polyphenols, and alpha-lipoic acid have shown beneficial effects in reducing oxidative stress and improving metabolic balance. Although antioxidants cannot completely cure diabetes, they may serve as valuable complementary strategies to limit complications and improve patient health.

Keywords: Type 2 diabetes, oxidative stress, reactive oxygen species, antioxidants, vitamin C, vitamin E, glutathione, polyphenols, alpha-lipoic acid, diabetic complications.

Introduction: Type 2 diabetes mellitus (T2DM) represents one of the most pressing health issues of the 21st century. According to the International Diabetes Federation (IDF), the global prevalence of diabetes is projected to reach 643 million by 2030 if current trends continue. While T2DM has traditionally been linked to insulin resistance, β -cell dysfunction, and impaired glucose metabolism, a growing body of evidence suggests that oxidative stress plays a pivotal role in both the initiation and progression of the disease.

Oxidative stress occurs when there is an imbalance between the excessive generation of reactive oxygen species (ROS) and the body's antioxidant defense systems. Elevated ROS levels are capable of damaging vital biomolecules including lipids, proteins, and nucleic acids. This cumulative cellular injury contributes not only to hyperglycemia-induced toxicity but also to the development of long-term diabetic complications.

Given the central role of oxidative stress in the pathophysiology of T2DM, antioxidant-based strategies have gained considerable attention. Researchers are increasingly investigating whether dietary or pharmacological antioxidants may help in preventing or slowing the progression of diabetes-related complications.

Biochemical Basis of Oxidative Stress in Type 2 Diabetes

Hyperglycemia and ROS Production

Chronic hyperglycemia is one of the primary drivers of ROS overproduction in diabetic patients. Several mechanisms are involved:

1. Glucose auto-oxidation: Glucose molecules in the bloodstream undergo spontaneous oxidation, generating superoxide anions and hydrogen peroxide.
2. Protein glycation: Persistent hyperglycemia leads to the non-enzymatic glycation of proteins, producing advanced glycation end-products (AGEs). These AGEs interact with receptors (RAGE), initiating further ROS generation and inflammatory responses.
3. Mitochondrial dysfunction: The overload of glucose in mitochondria increases electron leakage from the electron transport chain, which in turn accelerates superoxide formation.

Impact on Cellular Structures

The overproduction of ROS has widespread cellular consequences:

- Lipid peroxidation damages plasma and organelle membranes, disrupting cell integrity and signaling.
- Protein oxidation alters enzymatic activity, leading to reduced insulin receptor sensitivity.
- DNA damage results in mutations and may impair β -cell survival and proliferation.

These molecular disruptions amplify insulin resistance and exacerbate β -cell dysfunction, creating a vicious cycle that accelerates the progression of T2DM.

Antioxidant Defense Systems in the Human Body

Under physiological conditions, the human body has robust mechanisms to counteract oxidative stress. Key enzymatic antioxidants include:

- Superoxide dismutase (SOD) – converts superoxide anions into hydrogen peroxide.
- Catalase – breaks down hydrogen peroxide into water and oxygen.
- Glutathione peroxidase (GPx) – reduces lipid hydroperoxides and hydrogen peroxide using glutathione as a substrate.

In addition, non-enzymatic antioxidants such as vitamins C and E, uric acid, and polyphenolic compounds obtained from the diet provide an additional layer of protection. However, in individuals with T2DM, these defense systems are often insufficient to neutralize the elevated ROS load.

Therapeutic Potential of Antioxidants

Vitamins and Nutrients

Numerous clinical and experimental studies have demonstrated the beneficial effects of antioxidants in managing oxidative stress in T2DM:

1. Vitamin C (ascorbic acid): This water-soluble antioxidant directly scavenges ROS and regenerates vitamin E, enhancing overall antioxidant capacity. Supplementation has been shown to improve endothelial function and reduce oxidative DNA damage in diabetic patients.
2. Vitamin E (tocopherol): Being lipid-soluble, it primarily protects cell membranes from lipid peroxidation. Studies suggest that vitamin E supplementation improves insulin sensitivity and reduces markers of oxidative stress.
3. Glutathione: Known as the “master antioxidant,” glutathione plays a crucial role in detoxification and maintaining cellular redox balance. Diabetic patients often present with depleted glutathione levels.
4. Polyphenols: Found in green tea, grapes, and berries, these compounds have dual effects — direct antioxidant activity and enhancement of insulin sensitivity. Resveratrol, a well-studied polyphenol, has demonstrated protective effects against mitochondrial dysfunction.
5. Alpha-lipoic acid (ALA): Unique in being both water- and fat-soluble, ALA improves mitochondrial energy metabolism while reducing oxidative stress. It has been clinically used to alleviate diabetic neuropathy.

Pharmacological Advances

Beyond dietary supplements, pharmacological approaches are being developed to enhance antioxidant defense:

- Synthetic antioxidants such as N-acetylcysteine (NAC) increase intracellular glutathione levels.
- Mitochondria-targeted antioxidants like MitoQ are designed to specifically reduce ROS at the source of their production.

- Enzyme mimetics attempt to replicate the activity of SOD or catalase, thereby reducing oxidative stress in diabetic tissues.

Discussion: While antioxidant therapy holds great promise, several limitations must be acknowledged. First, clinical trials have produced mixed results, with some showing significant benefits while others report minimal effects. This variability may be attributed to differences in dosage, duration, bioavailability, and patient populations. Second, excessive antioxidant supplementation may paradoxically disrupt normal redox signaling, leading to unintended side effects. Thus, determining the optimal dose and treatment duration remains a critical challenge.

Nonetheless, combining antioxidants with standard antidiabetic therapies — including metformin, insulin sensitizers, and lifestyle interventions — may provide synergistic benefits. Metformin itself has mild antioxidant properties, which may explain its long-term protective effects beyond glucose control.

Conclusion: Oxidative stress is now recognized as a fundamental contributor to the pathogenesis and complications of type 2 diabetes. The overproduction of ROS disrupts normal cellular function, accelerates insulin resistance, and damages critical biomolecules. Antioxidants, both dietary and pharmacological, provide a promising avenue for mitigating these effects.

Future research must focus on refining antioxidant-based therapies through well-designed clinical trials, with particular attention to optimal dosing, bioavailability, and long-term safety. Ultimately, incorporating antioxidants into a holistic management plan may help reduce the burden of T2DM and improve patient outcomes worldwide.

References:

1. American Diabetes Association. (2023). Standards of Medical Care in Diabetes—2023. *Diabetes Care*, 46(Supplement 1), S1–S154.
2. Giacco, F., & Brownlee, M. (2010). Oxidative stress and diabetic complications. *Circulation Research*, 107(9), 1058–1070.
3. Rani, V., Deep, G., Singh, R. K., Palle, K., & Yadav, U. C. S. (2016). Oxidative stress and metabolic disorders: Pathogenesis and therapeutic strategies. *Life Sciences*, 148, 183–193.
4. Maritim, A. C., Sanders, R. A., & Watkins, J. B. (2003). Diabetes, oxidative stress, and antioxidants: A review. *Journal of Biochemical and Molecular Toxicology*, 17(1), 24–38.